

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

BIOREMEDIATION OF CONTAMINATED SOIL

A Thesis presented in partial fulfilment of the requirements for the degree
of Master of Technology in Environmental Engineering at Massey
University.

ROBERT JASON WOTHERSPOON

1997

ABSTRACT

The release of contaminants into the environment is inevitable. Contaminants are released through manufacture and use of products and as a result of treatment and disposal of wastes. Upon release to the environment, contaminants move and respond to a number of interrelated natural and man made factors.

Penta-chloro-phenol (PCP) is one such contaminant that has been released into the environment and is known to have serious long term environmental effects. The objective of this study was to determine the effectiveness of biological processes to remediate soil contaminated with Penta-chloro-phenol (PCP). This thesis reviews mechanisms by which soil is contaminated, processes available to remediate soils, and in particular, process requirements for successful bioremediation.

The abilities of bacteria to degrade PCP from soil contaminated with PCP was evaluated. Solid phase and slurry phase experiments were examined for their effect on PCP concentration over a four month period at the Department of Technology, Massey University. The objectives of this study were (1) To determine if aeration and inoculation of soil in-situ could produce significant removal of PCP. (2) Determine the effect of concentration on bioremediation rates. (3) Compare in-situ treatment with bio-slurry treatments.

The experiments showed that it is possible to remove up to 95 % of PCP from contaminated soil by inoculation with bacteria. Inoculum size and aeration were shown to be critical factors in affecting the rate of degradation. The larger the initial inoculum the greater the rate of degradation. Without aeration the inoculum was unable to significantly degrade PCP. The bio-slurry confirmed that PCP could be removed readily from soil to an aqueous state. In an aqueous state PCP is degraded at a faster rate than when it is incorporated into the soil matrix.

The results of this work is to show that soil rehabilitation by way of biodegradation is a feasible and attractive process.

ABSTRACT

1. INTRODUCTION	1
2. CHEMICALS IN SOIL.....	2
2.1 INTRODUCTION	2
2.2 PHYSICAL PROCESSES CONTROLLING THE TRANSPORT OF CONTAMINANTS IN THE AQUEOUS PHASE.....	2
2.3 TRANSPORT OF NON-AQUEOUS PHASE LIQUIDS IN THE SUB SURFACE	6
3. REMEDIATION TECHNIQUES	18
3.1 INTRODUCTION	18
3.2 PHYSICAL TECHNIQUES.....	19
3.3 CHEMICAL TECHNIQUES	23
3.4 THERMAL TECHNIQUES.....	25
3.5 SOLIDIFICATION AND STABILISATION	26
3.6 BIOLOGICAL TECHNIQUES.....	27
3.7 REMEDY SELECTION.....	29
4. BIOREMEDIATION OF PCP CONTAMINATED SOIL	30
4.1 INTRODUCTION	30
4.2 MICROBIOLOGICAL PRINCIPLES INFLUENCING BIOREMEDIATION.....	32
4.3 ABIOTIC TRANSFORMATION REACTIONS	33
4.4 BARRIERS TO BIODEGRADATION.....	37
4.5 IN SITU BIOREMEDIATION	49
4.6 SLURRY TREATMENT	54
4.7 CONCLUDING REMARKS.....	58
5. MATERIALS AND METHODS	60
5.1 SOIL	60
5.2 BACTERIA	61
5.3 MEDIA	62
5.4 CHEMICALS	62
5.5 AERATION EQUIPMENT.....	62
5.6 GLASSWARE.....	64
5.7 SLURRY REACTOR	64
6. ANALYTICAL PROCEDURES	65
6.1 SOIL pH.....	65
6.2 SOIL MOISTURE.....	65
6.3 TOTAL NITROGEN.....	65
6.4 PCP EXTRACTION	66
6.5 CHLORIDE EXTRACTION.....	68
6.6 MICROBIOLOGICAL ANALYSIS	68
6.7 ISOLATION MEDIA.....	69
6.8 STATISTICAL ANALYSIS OF DATA	69
7. EXPERIMENTAL PROCEDURE.....	70
7.1 EXPERIMENT ONE.....	70
8. EXPERIMENT 2	72
9. EXPERIMENT 3	74
10. RESULTS AND DISCUSSION	76
10.1 PRELIMINARY REMARKS.....	76
10.2 EXPERIMENT 1: RESULTS.....	77

10.3 DISCUSSION EXPERIMENT 1	82
10.4 EXPERIMENT 2.....	85
10.5 RESULTS.....	85
10.6 FIGURES 1.7-1.12	87
10.7 DISCUSSION EXPERIMENT 2	90
11. EXPERIMENT 3.....	91
11.1 RESULTS.....	92
11.2 DISCUSSION EXPERIMENT 3	92
12. CONCLUSIONS.....	96

LIST OF TABLES

Table 1 Critical Height for Perchloroethylene to Penetrate Water Saturated Media	14
Table 2 Mechanisms influencing the fate of contaminants in environments	32
Table 3 Selected hydrolysis half lives for a variety of organic compounds	34
Table 4 Microbial Growth response to Phenyldecanes	42
Table 5 Adaptation period (days)	46
Table 6 Soil Characteristics	60
Table 7 Cation Exchange Properties of soil	60
Table 8 Chemostat operational conditions	61
Table 9 Composition of Feed	62
Table 10 Vertical extent of PCP degradation.	78
Table 11 PCP half lives	83
Table 12 PCP degrading bacteria concentrations	86
Table 13 Residual PCP concentration	91
Table 14 Comparison of PCP half life in solid and slurry phase	94

LIST OF FIGURES

Figure 1 Relative permeability versus water saturation	8
Figure 2 LNAPL distribution after spillage	10
Figure 3 Depression of capillary fringe and water table	10
Figure 4 Rebounding of water table	11
Figure 5 Distribution of DNAPL after small volume spilled	15
Figure 6 Distribution of DNAPL after moderate volume spilled	16
Figure 7 Distribution of DNAPL after large volume spilled	16
Figure 8 Pathway of carbaryl degradation	40
Figure 9 Lag Adaptation periods.	44
Figure 10 Effect of chemical structure on adaptation periods.	46
Figure 11 Effect of substrate concentration on length of adaptation period	47
Figure 12 Differences in the adaptation period as a function of substrate concentration.	47
Figure 13 Slurry phase bioreactor	57
Figure 14 Schematic of chemostat	61
Figure 15 Experimental layout	63
Figure 16 Column 1 Experiment 1, PCP 130 mg/kg	79
Figure 17 Column 2 Experiment 1, PCP 92 mg/kg	79
Figure 18 Column 3 Experiment 1, PCP 50 mg/kg	80
Figure 19 Column 4 Experiment 1, PCP 45 mg/kg	80
Figure 20 Column 5 Experiment 1, PCP 48 mg/kg no aeration	81
Figure 21 Column 6 Experiment 1, PCP 51 mg/kg no inoculum	81
Figure 22 Column 1 Experiment 2, 1×10^7 cells	87
Figure 23 Column 2 Experiment 2, 1×10^6 cells	87
Figure 24 Column 3 Experiment 2, 1×10^5 cells	88
Figure 25 Column 4 Experiment 2, 1×10^7 cells	88
Figure 26 Column 5 Experiment 2, 1×10^5 cells	89
Figure 27 Column 4 Experiment 2, 1×10^4 cells	89
Figure 28 PCP concentration in Soil and aqueous phase	93
Figure 29 Disappearance of PCP in bioslurry reactor	93

REFERENCES

APPENDICES

ACKNOWLEDGMENTS

This completion of this thesis involved assistance from many people. I am grateful to all those who helped, and particularly wish to thank the following people.

Professor Rao Bhamidimarri is especially thanked for his supervision of the entire project. His willingness to give time and expert assistance whenever needed was greatly appreciated.

I wish to thank Dr Bob Chong and Dr Graeme Manderson whose expert advice and time throughout the duration of the project was greatly appreciated.

Mr John Alger is thanked for his construction of experimental equipment and for his technical assistance throughout the project.

The trustees of the A.D.B. Williams Scholarship are thanked for the provision of funds for this project.

I am especially grateful to my parents, and wish to thank them for all their support and encouragement throughout my education, especially during my years at University.

1. INTRODUCTION

The total land area on Earth is approximately 14500 million hectares of which 13250 million hectares is ice free. Land use patterns are mainly determined by interactions between climate, geography, geology, human and economic processes. Land can be categorised into four categories: arable and permanent cropland, permanent pasture, forests and land for general use including land unused in urban areas, and waste and barren land. The latter encompasses a third of the earth's surface, of which more than 3,000 million hectares is suspected to be exposed to chemicals¹.

All categories of land are open to pollution and consequently are hazardous to the environment, attention has been focussed on land categorised under general use because this represents a more direct risk to human health and thus receives the greater publicity in the case of incidents.

Contaminated land results from a wide range of human activities including industrial discharge processes and the disposal of waste. New Zealand has an advantage over old world countries such as the United Kingdom and United States of America in that there is not a legacy of nearly two centuries of industrial operation. Despite this New Zealand has sites which are as heavily contaminated as some overseas industrial sites. Remedial technique experience with respect to contaminated land is brief, dating back only to the 1970's. It is important in the long term to develop new and effective means of soil decontamination.

Bioremediation of soils is a novel technique that is slowly gaining acceptance despite having significant cost advantages over other remediation technologies. The main reason for this lack of acceptance being a reluctance on the part of engineers to utilise a treatment option that was generated outside their field.

The aim of this thesis is to add to this body of knowledge